



# FERTI-ZINC 10%

## MICRONUTRIENT COMPLEX

**GENERAL USE INFORMATION – READ ENTIRE LABEL BEFORE USING THIS PRODUCT**

### GENERAL INFORMATION

FERTI-ZINC 10% is recommended for field, row vegetable, fruit, tree, vine, ornamental, and turf crops. FERTI-ZINC 10%, alone or with pesticides or fertilizers, may be applied directly to soil or to the foliage of growing crops. FERTI-ZINC 10% may be applied via ground sprayers (high volume or low-volume), aircraft or drip irrigation systems.

### GENERAL DIRECTIONS FOR USE

**CONSULT YOUR LOCAL FERTIZONA REPRESENTATIVE OR OTHER AGRICULTURAL SPECIALIST FOR SPECIFIC CROP RECOMMENDATIONS.**

Soil application of FERTI-ZINC 10% is recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliar-applied FERTI-ZINC 10% most efficiently. Immediate post-irrigation or post-rainfall applications give best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Biuret Urea per 100 gals. of spray solution) will enhance foliar uptake of FERTI-ZINC 10%.

AVOID foliar applications of FERTI-ZINC 10% when plants are under moisture stress.

AVOID application to deciduous tree crops during bloom.

COMPATIBILITY: FERTI-ZINC 10% is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

MIXING: ALWAYS add FERTI-ZINC 10% to spray tank BEFORE adding pesticides.

### NOTICE

**READ ENTIRE LABEL BEFORE USE.**

**IF TERMS ARE UNACCEPTABLE RETURN AT ONCE – UNOPENED.**

Fertilizer Company of Arizona, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Fertilizer Company of Arizona, Inc. In no case shall Fertilizer Company of Arizona, Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. The buyer shall assume all such risks. Fertilizer Company of Arizona, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Distributed By:

**FERTILIZER COMPANY OF ARIZONA, INC.**

2850 S. Peart Road • Casa Grande, Arizona 85193-9024 • (520) 836-7477

### GUARANTEED ANALYSIS

Zinc (Zn) ..... 10.0%  
Sulfur (S) ..... 4.0%  
Derived from zinc sulfate and zinc lignin sulfonate.

*MSDS available upon request.*

### NET WEIGHT:

**11 LBS. PER GAL. AT 68° F.**

5 Gals. (18.925 L), 55.0 lbs. (24.948 kg)

### STORAGE AND DISPOSAL

Do not use aluminum or copper (including brass or bronze) components in any handling or storage equipment for this product. Mild steel components should not be used except for emergency, or temporary, short-term service. Use stainless steel (316 or better), polyolefin (polyethylene, polypropylene, etc.), fiberglass, or other high-quality acid-resistant plastic materials. Check with equipment manufacturer for compatibility of components prior to use with this product.

STORAGE INSTRUCTIONS: Protect from freezing. Store above 40°F. Freezing will not harm product's effectiveness. Store in original container tightly closed, do not allow water to be introduced into contents of this container.

CONTAINER DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Triple rinse (or equivalent), then offer container for recycling, or puncture and dispose in a sanitary landfill or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### SHAKE, STIR OR SWIRL CONTENTS BEFORE USING

### SPECIFIC DIRECTIONS FOR USE

ALL CROPS:\*

#### SOIL APPLICATIONS

Soil Fertility Maintenance ..... 1-2 qts. per acre  
Light Deficiency ..... 2-4 qts. per acre  
Severe Deficiency ..... 1-3 gals. per acre

\*The lower rates are for field, row, and vegetable crops. The higher rates are for fruit, tree, vine, ornamental, and turf crops.

#### FOLIAR APPLICATIONS

For rates (above) higher than 2 qts. per acre, split applications of 2 qts. or less each are preferable. Applications should be separated by a minimum of 10-14 days.

#### DILUTE APPLICATIONS

Apply the suggested dose of FERTI-ZINC 10% in enough water for good coverage.

#### CONCENTRATE APPLICATIONS

Use up to ½ gallon of FERTI-ZINC 10% in a minimum of 25 gals. of water per acre.

#### AERIAL APPLICATIONS

Up to 1 qt. of FERTI-ZINC 10% per 5 gals. of spray may be used, but more dilute solutions will improve coverage and results.

#### DRIP IRRIGATION APPLICATIONS

Apply the recommended amount of product in split application 1-2 weeks apart throughout the growing season.

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**